



Mid-late Holocene climate variability in the Indian monsoon: Evidence from continental shelf sediments adjacent to Rushikulya river, eastern India



Yadav Ankit ^a, Prem Kumar ^a, Ambili Anoop ^a, Praveen K. Mishra ^b, Saju Varghese ^{c,*}

^a Indian Institute of Science Education and Research Mohali, Manauli, Punjab, 140306, India

^b Wadia Institute of Himalayan Geology, Dehradun, 248001, Uttarakhand, India

^c Geological Survey of India, Marine and Coastal Survey Division, Salt Lake, Kolkata, 70091, India

ARTICLE INFO

Article history:

Received 20 September 2016

Received in revised form

12 December 2016

Accepted 21 December 2016

Available online 1 March 2017

Keywords:

Indian monsoon

Eastern India

Continental shelf

Geochemistry

Holocene

ABSTRACT

We present elemental and grain-size distributions obtained from the sediment core of the continental shelf adjacent to the Rushikulya river mouth, eastern India to quantify the paleoclimatic changes. The retrieved 1.60 m long well dated core spans the past ca. 6800 cal BP. The modern spatial distribution of grain size and geochemistry of the inner-mid shelf sediments has been carried out to understand the seafloor morphology and sedimentary processes. Based on the modern investigations, the proportion of particle size (clay vs sand) and variation in elemental values (TiO₂ vs Al₂O₃) has been used to interpret the changes in terrigenous supply. The grain-size and elemental distribution data from the core sediments indicates a period of enhanced surface water runoff from 6800 to 3100 cal BP followed by a drier condition (3100 cal BP to present) suggesting weakening of monsoon. The weakening of the monsoonal strength is coeval with other records from the Indian sub-continent and suggests response of Indian monsoon to changing solar insolation during late Holocene.

© 2016 Elsevier Ltd and INQUA. All rights reserved.

1. Introduction

The understanding of the past and present climatic variability in the Indian monsoon realm is critical to produce better future climate forecast models. However, the available long term continental palaeomonsoon record has limited geographical coverage with majority of research carried out in NW India (e.g., Prasad et al., 1997; Enzel et al., 1999; Prasad and Enzel, 2006; Prasad et al., 2014a; Raj et al., 2015; Amekawa et al., 2016) and Himalayan region (Phadtare, 2000; Demske et al., 2009; Wünnemann et al., 2010; Anoop et al., 2013a; Leipe et al., 2014; Rawat et al., 2015; Mishra et al., 2015; Kotlia et al., 2015). To understand the variability of monsoon rainfall and the physical mechanisms inherit in it, a large dataset covering longer time scales is required from various climate sensitive regimes across the Indian subcontinent. Barring a few exceptions (e.g., Khandelwal and Gupta, 1999; Khandelwal et al., 2008) limited to palynological investigations

on lacustrine sediments, the Holocene climate changes are yet to be evaluated from eastern India.

The sedimentary bodies on continental shelves have been widely used to infer the sea level fluctuations (Milliman and Emery, 1968; Gingele et al., 2004; Zhao et al., 2008; Yoo et al., 2014) and variations in terrigenous sediment input (Mendes et al., 2010; Nizou et al., 2010; Zheng et al., 2010; Rosa et al., 2011; Wang et al., 2014; Perez et al., 2016; Tu et al., 2016) interpreted in terms of past environmental changes. The terrigenous coarse- and fine-grained sediments supply to the continental shelf is mostly controlled by fluvial discharge (Prins et al., 2000; Briceño-Zuluaga et al., 2016). The paleoclimate records from continental shelves provide critical information connecting continental with deep marine records (González-Álvarez et al., 2005). However, paleoclimate records from continental shelf areas of the Indian subcontinent are very scarce. In this study, we present a continuous record of the mid-late Holocene climate obtained from a 1.60 m long sediment core (VC-04) retrieved from the continental shelf sediments adjacent to the Rushikulya river mouth (Fig. 1). The understanding of modern sedimentation patterns in the continental shelf has been used to refine palaeoenvironmental interpretations and gain a long term perspective on environmental variability. The

* Corresponding author. Geological survey of India, Marine and Coastal survey division, Block DK 6, Sector II, Salt Lake, Kolkata, 70091, India.

E-mail address: saajuvarghese@gmail.com (S. Varghese).




Fig. 1. Map showing study area and location of the retrieved core (VC-04) for paleoclimate reconstruction. The blue, yellow and red circles denote the location of vibro, gravity and grab samples used to understand the modern continental shelf dynamics. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

elemental and grain-size distributions supplemented with sedimentation rate calculations obtained from the sediment core are used to quantify the paleoclimatic changes.

The objectives of this present study are: i) to identify suitable proxies for paleoenvironment reconstruction by establishing the linkages between contemporary processes and sediment distribution in the modern Rushikulya river mouth and adjacent continental shelf sediments; ii) use of climate sensitive proxies to reconstruct paleoenvironmental conditions during mid-late Holocene; and iii) regional comparison of the climate data to understand the physical mechanisms that control the spatial temporal variability of monsoon precipitation in the region.

2. Study area

2.1. Regional setting

The investigated region includes inner to mid continental shelf of Rushikulya river mouth extends between $19^{\circ}11.9'$ – $19^{\circ}23.4'$ N and $85^{\circ}00.6'$ – $85^{\circ}13.9'$ E covering an area of 245 km². The Rushikulya river emerges from the slopes of Araha bity and Kutror hills of Eastern ghats with a drainage area of 82,100 km² running for approximately 165 km before meeting the Bay of Bengal at Puruna Bandha of Chhatarpur region (Jain et al., 2007). The average annual

water discharge of the Rushikulya river is ~1800 million m³ with discharge highly seasonal due to the monsoonal conditions (Panigrahy et al., 1999).

2.2. Regional climate and geomorphology

The climate over the east coast of India is mainly tropical in nature administered by the SW monsoon between June and September, and NE monsoon during October to December. The study area receives an average annual rainfall of ~1200 mm with SW monsoon accounts for ~80% of the total precipitation (Jain et al., 2007). The maximum and minimum temperatures recorded in the region are 45°C and 12°C respectively (Jain et al., 2007). The wind force over the region is fairly high during the period of south-western monsoon and the whole of Odisha coast is vulnerable to frequent cyclonic storms during October–November (Mascarenhas, 2004).

The high energy waves from the south and the southeast during the monsoon season generate the long shore current that heads towards north and is reversed during winter months due to prevailing northeasterly waves generating littoral current that runs towards south (Mishra et al., 2001). The coastal region of the study area is characterised by major geomorphic features like beach ridges, sand spits, and barrier spits (Markose et al., 2016). The

coastline adjoining the study area is straight trending 40°–50° (Varghese et al., 2015). The continental shelf in this area is smooth and gently sloping as inferred from the smooth curves of the isobaths running more or less parallel to the coastline (Varghese et al., 2015).

2.3. Regional geology

The mineral assemblages over the study region are derived from Eastern Ghats Mobile belt (EGMB) (Bhattacharya and Sengupta, 1994). The litho units of EGMB are divided longitudinally into four major lithozones comprising of eastern khondalite zone (EKZ), charnockite migmatite zone (CKZ), western khondalite zone (WKZ) and western charnockite zone (WCZ) (Ramakrishnan et al., 1998). The sediments deposition over beach of southern coast Odisha, Chhatarpur and Rushikulya area is known for its mineral reserves. The minerals which are economically important present in the region include ilmenite, garnet, sillimanite and rutile (Rao et al., 2001). The coastal charnockite provide main sources for garnet as beach placer, whereas garnet-quartz and sillimanite garnet quartz schists serve as important sources for rutile (Rao and Misra, 2009). These minerals over study area are carried out in form of detrital grains by seasonal (monsoon) rivers and further redistributed by marine long shore current (Rao, 1957).

3. Methodology

3.1. Sampling

The sediments cores from the inner-mid continental shelf adjacent to the Rushikulya river mouth were retrieved using 5 m long Nino's type gravity and vibro corer from the desired locations (Fig. 1) during the R.V Samudra Kaustubh cruise ST-223 in 2012–2013. The uppermost 0–5 cm of vibro and gravity cores has been used to understand the modern sedimentation processes along the Rushikulya river continental shelf. Additionally, grab samples also have been collected using medium sized Van Veen/Peterson grab with an approximate penetration up to 25 cm. All the gravity and vibro core samples collected in PVC liner were split and logged visually as per variation in sediment colour, texture, compaction and relative proportion of biogenic and terrigenous constituents. Immediately after collection, samples were kept frozen till brought to lab for further studies.

The down core variation of major oxides and grain size distribution is studied for the ca. 160 cm long vibro core VC-04 collected (42.8 m water depth) adjacent to the Rushikulya river mouth. The elemental and grain size measurements were carried out at 10 cm resolution.

3.2. Analytical methods

3.2.1. Geochemistry

The sediment samples for XRD analyses were crushed into a fine grained powder using a diamond mortar followed by wet grinding. The pestle was put on a glass sample holder that was placed on the sample rotation stage (120 rpm) of the instrument. The analysis were carried out on a Rigaku Ultima IV diffractometer equipped with a 3 kw sealed tube Cu K α X-ray radiation (generator power settings, 40 kV and 40 mA) and a DTex Ultra detector using parallel beam geometry. The data were collected over an angle range of 5°–90° with a scanning speed of 5° per minute with a 0.02° step.

For elemental analysis, approximate 20 gm of air dried sub-samples were desalinated thrice using distilled water before chemical treatment. The desalinated samples were crushed in an agate mortar until entire quantity passed through 100–120 ASTM

mesh size. The powdered samples were digested with a mixture of concentrated HF–HNO₃–HClO₄ following the method of Zhang and Liu, 2002. A fraction (~0.25 gm) from each sub-sample was treated with aqua regia followed by treatment with Hydrofluoric acid (HF) to remove the silicates. Perchloric acid was used to remove the organic content in the sediments before the samples were made up to a 100 ml solution using the dilute nitric acid. The samples in the nitric medium fed to a Varian 720-ES Inductively Coupled Plasma Optical Emission Spectrometer (ICP-OES) for major oxide analysis. P₂O₅ is determined by color development with Ammonium Molybdate-Ammonium-Meta-Vanadate solutions with subsequent measurement on Spectrophotometer at 465 nm. Precision for major oxides is generally better than 2%.

3.2.2. Grain size analysis

The sample pre-treatment for grain-size analysis includes organic matter removal by addition of H₂O₂ followed by dispersal of aggregates using ammonia. The sediment samples were then washed with distilled water and oven-dried at 45 °C. In the present case, grain size distribution (GSD) of the samples was studied using a combination of sieve method for sand dominated sediments and pipette analysis for silt and clay rich sediments (Folk, 1980; McCave and Syvitski, 1991; Miall, 2006).

3.2.3. Chronology

The age model for VC-04 is obtained based on the four AMS (accelerator mass spectrometry) ¹⁴C dates of bulk organic sediments (Table 1). The ¹⁴C dating points were chosen according to changes in the lithology and elemental values. Radiocarbon dating was carried out at Radiochronology lab in University of Laval, Canada. The obtained ¹⁴C dates were then converted from conventional ¹⁴C ages into the calendar ages using OxCal 4.1 software (Ramsey, 2001, 2009) with the IntCal 13 calibration curve (Reimer et al., 2013). The age-depth relationship was modeled using linear interpolation between adjacent radiocarbon dates.

3.3. Data analysis

Principal component analysis (PCA) was carried out for geochemical concentrations of the surface and core sediments to explore element associations and their origins. PCA analysis is a multivariate technique which explains the major variations within dataset by generating factors which represent cluster of interrelated variables (Reid and Spencer, 2009). The analysis enables a reduction in the dimensionality of a dataset by means of small number of components (Loska and Wiechula, 2003). The principal components (PC) should account for ~75% of the total variance (Morrison, 1967). The varimax rotations of the generated PCs are performed to maximize the variation in the data for easy interpretation (Juahir et al., 2011; Sappa et al., 2014). However, the obtained PCs of the geochemical data are readily interpreted; therefore, varimax rotation has not been applied. PCA analysis was performed using XLSTAT, an add-in software package for Microsoft Excel (Addinsoft Corp.).

Table 1
AMS radiocarbon dates for the core VC-04 derived from bulk organic matter.

No	Depth (cm)	Lab no.	Age (¹⁴ C)	Age (cal BP) (1 σ)
RS-5	0	ULA-6259	112.73 ± 0.18*	Modern
RS-6	15	ULA-6260	880 ± 15	788 ± 43
RS-7	65	ULA-6261	2975 ± 15	3139 ± 51
RS-8	160	ULA-6262	5980 ± 20	6816 ± 36

* Denotes value in pMC (percent modern carbon).

4. Results

4.1. Modern data

4.1.1. Geochemistry and grain size analysis of surface sediments

The XRD investigation of the modern sediments revealed that the dominant siliciclastic minerals constitute quartz, biotite, orthoclase, rutile, sillimanite, wollastonite and muscovite. The aragonite and kaolinite represent dominant carbonate and clay mineral phases respectively.

The PCA analysis of the surface sediments extracts two principle components axis explaining ~79.68% of the total variance (F1-55.51%; F2- 24.17%) (Fig. 2a). The important characteristic of the




Fig. 2. Biplot of F1 axis vs. F2 axis based on the principal component analysis of geochemical parameters from (a) continental shelf surface sediments and (b) core sediments (VC-04).

PCA analysis is the existence of two clusters for the 'terrigenous' elements that are anti-correlated (Fig. 2a) (Table 2). The F1 shows high correlation with the elements Fe₂O₃, Al₂O₃, K₂O and Na₂O, whereas F2 suggests its close relation with the elements TiO₂ and MnO (Fig. 2a). The elemental concentrations of CaO shows low positive loadings in F1 and F2 axis attributed to the contribution from dual sources- calcium bearing detrital minerals and authigenic aragonite shells present within the surface sediments. Based on the PCA results, Al₂O₃ and TiO₂ have been taken as representative elements to understand the modern dynamics in the inner to mid continental shelf. The trend in the elemental distribution of Al₂O₃ display high values towards the deeper part of the continental shelf, whereas TiO₂ shows higher values close to the shore region (Fig. 3b and 3c).

The grain size exhibit specific variability trends in their inner-mid shelf distribution along the study area (Fig. 3d–f). The grain size distribution of the shelf sediments shows high variability characterised by ~2–97% of sand, ~1–93% of silt and ~1–21% of clay. The spatial distribution of the grain size in the surface sediments shows an increasing trend of finer fractions (e.g. silt and clay) with increasing water depth, whereas coarser fraction (e.g. sand) shows a decreasing trend (Fig. 3d–f).

4.2. Core data (VC-04) from continental shelves

4.2.1. Lithology

Based on the geochemistry and visual observation, lithology of the core sediments (VC-04) shows two major distinct units. The unit I (0–0.72 m) comprises of olive gray loose silty clay with forams, whereas unit II (0.72–1.60 m) consists of very fine sand with silt and frequent shell fragments.

4.2.2. Chronology

The uncertainty associated with ¹⁴C ages of the bulk organic matter (OM) derived from the near shore marine environment involves incorporation of old plant-derived terrigenous organic carbon resulting in anomalously old ¹⁴C ages and age reversals (Eglinton et al., 1997; Zheng et al., 2002). However, the modern age of the bulk OM from the top of the sediment cores provides evidence for absence of possible contamination. Additionally, stratigraphically consistent ¹⁴C dates further indicate the reliability of the ¹⁴C dates (Table 1). The sedimentation rates of the VC-04 displays elevated sedimentation rate (0.026 cm/yr) in the lower part (160–70 cm) of the core, whereas in the upper part (70–0 cm) the sedimentation rate was low and stable (average = 0.020 cm/yr) (Fig. 4).

4.2.3. Geochemical and grain size data of VC-04


The principal component analysis of the core sediments (VC-04) extracted two principal components that account for 92.98% of the total variance. The first axis (F1) explains 77.21% of the total variance, whereas the second PCA axis (F2) explains 15.77% of the total variance (Fig. 2b). The first axis (F1) shows high positive factor loadings for Al₂O₃-Fe₂O₃-K₂O-Na₂O, whereas the second axis (F2) shows high positive factors loadings for TiO₂. The PCA analysis of core sediment (VC-04) reveals similarities with the modern shelf distribution showing anti-correlation of TiO₂ with other detrital elements (Al₂O₃-Fe₂O₃-K₂O-Na₂O). The Al₂O₃ and TiO₂ of the core sediments range from 4.06 to 16.15% and 0.46–1.81% respectively. During 6800 to 3100 cal BP the core shows a low concentration of Al₂O₃, whereas the upper part (3100 cal BP to present) display high and stable contribution of Al₂O₃ (Fig. 4). The core VC-04 shows a rather high TiO₂ from 6800 to 3100 cal BP, and relatively low values from 3100 cal BP to present (Fig. 4).

The particle size data for the core sediments shows fluctuations

Table 2

Pearson correlation coefficient matrix (significant correlations denoted in bold) for the geochemical parameters of the inner-mid shelf surface sediments.

Variables	Al ₂ O ₃	Fe ₂ O ₃	MgO	CaO	MnO	Na ₂ O	K ₂ O	TiO ₂	P ₂ O ₅
Al ₂ O ₃	1	0.892	0.997	0.052	−0.035	0.537	0.882	−0.272	0.953
Fe ₂ O ₃	0.892	1	0.896	0.151	0.403	0.404	0.838	0.189	0.969
MgO	0.997	0.896	1	0.059	−0.023	0.554	0.888	−0.259	0.953
CaO	0.052	0.151	0.059	1	0.018	−0.223	0.102	0.198	0.162
MnO	−0.035	0.403	−0.023	0.018	1	−0.078	0.070	0.935	0.210
Na ₂ O	0.537	0.404	0.554	−0.223	−0.078	1	0.594	−0.261	0.408
K ₂ O	0.882	0.838	0.888	0.102	0.070	0.594	1	−0.114	0.845
TiO ₂	−0.272	0.189	−0.259	0.198	0.935	−0.261	−0.114	1	−0.008
P ₂ O ₅	0.953	0.969	0.953	0.162	0.210	0.408	0.845	−0.008	1

**Fig. 3.** a) Bathymetry of the study area, and spatial distribution of b) Al₂O₃ (wt%), c) TiO₂ (wt%), d) Sand (%), e) Silt (%) and f) Clay (%) of the inner-mid continental shelf sediments adjacent to Rushikulya river mouth.

of ~5–96%, ~4–57% and ~0–46% for sand, silt and clay respectively. The grain-size data of the VC-04 shows two phases of grain size distribution, coarser sand fraction (average = 94.5%) between 6800 and 3100 cal BP and finer clay and silt sediments (average = 92.5%) from 3100 cal BP to present (Fig. 4).

5. Discussion

5.1. Modern proxy development

5.1.1. Grain size distribution

The sediment distribution patterns in the continental shelves are mostly controlled by type, proximity of the sediment sources and prevailing coastal conditions (e.g., Rosa et al., 2011). Grain size is mostly dependent on the process of weathering, erosion, transport, and sedimentation rate (Hjulstrom, 1935, 1939; Visher, 1969). In the study area, the modern sedimentation pattern follows a classical nature of sedimentation i.e. coarser size sediments deposited near shore reflecting high energy flow, while the finer

sediments toward the pelagic depth reflecting low energy flow.

5.1.2. Geochemical characterisation of the surface sediments

The relation between TiO₂ and Al₂O₃ are used to investigate the variation in detrital input, rainfall intensity and runoff contribution from the catchment (Minyuk et al., 2013), assuming that the Ti and Al both are conservative in nature during the weathering process (Boës et al., 2011). However, in the sediment from the Rushikulya River, there is poor correlation between Al₂O₃ and TiO₂ ($r = -0.27$) suggesting that both are either controlled by (i) grain size variability or (ii) having a different source of sediments. Fig. 3b, e and f shows that the Al₂O₃ concentration is highest at the deepest part of the study area following the trend of silt and clay size grain distribution. However, TiO₂ is dominant at the shallow part of the shelf and showing high correlation with sand suggesting its association with the coarser grain size sediments.

The surface sediments from the continental shelf show a high correlation between Al₂O₃ vs silt ($r = 0.60$), K₂O ($r = 0.89$), Fe₂O₃ ($r = 0.89$) and MgO ($r = 1.00$). The close association between these




Fig. 4. The temporal variation in sedimentation rate, geochemical (Al_2O_3 and $\text{TiO}_2\%$), and grain size (sand, silt and clay %) data of core VC-04.

elements may suggest the contribution of mica group of minerals. In addition, the geology of the catchment shows dominance of granitic gneiss and augen-gneiss which act as a provenance of the sediments rich in mica group of mineral. Furthermore, Ti is a common lithophile element that occurs in several minerals such as ilmenite (FeTiO_3), rutile, brookite (both TiO_2), and sphene (CaTi-SiO_5). The poor correlation of Ti with Fe and Ca suggests that rutile could be the key source of Ti in Rushikulya river sediments. The semi quantitative mineralogical analysis of the sediments also supported the presence of rutile in the sediments. The high concentration of TiO_2 near the river mouth following the trend of sand size sediments indicates that the grain size controls the elemental distribution of TiO_2 .

5.2. Interpretation of core sediment (VC-04)

The sharp lithological contrasts and the trends of geochemical and sedimentological variables indicate two distinct intervals with different environmental conditions in the study region during the last 6800 cal BP. The sandy nature of the lower half of the core near the Rushikulya river indicates the contribution of coarse grained fluvial sediment. In order to deposit sand layers on a high stand mid continental shelf, high energy processes must be invoked. The intermittent layers of sand and silt are due to occasional supply of large amount of sediment by Rushikulya river during monsoon season followed by period of less sediment supply. We exclude the influence of sea level fluctuations for the observed variations as investigations from western India continental margin shows minor sea level fluctuations for the past 7000 years (Hashimi et al., 1995). The sea-level curve presented from the eastern coast of southern India also indicates a gradual and slow sea-level rise since 9000 to 1800 yrs BP (Hameed et al., 2006). Additionally, the geochemical

and palynological analyses of relict mudflat from southern Saurashtra coast also reveals wet climatic conditions with simultaneous occurrence of marginally high sea-level between 4710 and 2825 cal BP (Banerji et al., 2015). The sharp sedimentary change observed at 3100 cal BP represents the transition from a shelf dominated by monsoonal terrigenous supply to a low-energy environmental condition.

5.3. Regional comparison of paleoclimate data

Grain-size and elemental distribution data from the core sediments (VC-04) indicates a period of enhanced surface water runoff from 6800 to 3100 cal BP followed by drier conditions to present (Fig. 5a). The enhanced precipitation in Rushikulya during ca. 6800–3100 cal BP is synchronous with the climatic reconstruction based on lake sediments from Nal Sarovar, central Gujarat (Prasad and Enzel, 2006) with high lake stand during ca. 4800–3000 cal BP (Fig. 5b). Likewise, the pollen and phytolith based investigations from western India, Pariyaj lake (Raj et al., 2015) and Wadhvana Lake (Prasad et al., 2014a) also shows a wet phase between ~4680 and 3500 cal BP 7500–5560 cal BP respectively. The wetter conditions in Rushikulya region during 6800–3100 cal BP is also in good agreement with the record based on carbon isotopes of sedimentary leaf waxes from core monsoon zone (Ponton et al., 2012) and mineralogical and isotopic investigation of Lonar core sediments from central India (Anoop et al., 2013b; Prasad et al., 2014b) (Fig. 5c and 5d). This humid phase also matches with the palaeorecords from mid-late Holocene from the ISM dominated Himalayan and Tibetan region (Dodia et al., 1984; Demske et al., 2009; Trivedi and Chauhan, 2009; Günther et al., 2015; Kotlia, 2015; Rawat et al., 2015; Mishra et al., 2015) (Fig. 5e) suggesting common forcing factor in governing the climate variability in the




Fig. 5. Regional comparison of VC-04 climate data with other paleoclimatic records: (a) Al_2O_3 (wt%) of the core sediment (present study); (b) schematic curve of climate change derived from Nal Sarovar data (Prasad and Enzel, 2006); (c) climate information from Lonar lake based on the presence of evaporite minerals (Anoop et al., 2013b; Prasad et al., 2014b); (d) carbon isotope data from biomarkers derived from the Godavari catchment (Ponton et al., 2012); (e) XRF profile of calcium (in cps) of Tso Moriri core sediment (Mishra et al., 2015); (f) climate information based on geochemical and palynological analyses of relict mudflat from Saurashtra coast (Banerji et al., 2015).

Indian sub-continent.

The onset of weakening of the ISM recorded from VC-04 is in phase with the gradual onset of aridity and lowering of the sea-level observed between 2825 and 1835 cal BP from the Saurashtra coast in western India (Banerji et al., 2015) (Fig. 5f). Furthermore, the results are also in line with the high resolutions record from NW India: Nal Sarovar lake (Prasad et al., 1997); Pariyaj Lake (Raj et al., 2015) Lunkaransar and Didwana Lakes (Wasson et al., 1984; Enzel et al., 1999). The findings of drier conditions in

Rushikulya region after ca. 3100 cal BP is good accordance with the sedimentary leaf waxes record from core monsoon zone (Ponton et al., 2012) and mineralogical and isotopic data from Lonar sediments in central India (Anoop et al., 2013b). The variations of the mid-late Holocene hydrological changes have shown significant linear response to gradually changing insolation (e.g., Prasad et al., 2014b; Mishra et al., 2015). The spatial variability in palaeodata has been mostly explained by changes in moisture sources or monsoonal teleconnections (e.g., ENSO, Indian ocean dipole (IOD)

(Hong et al., 2005; Sinha et al., 2011; Prasad et al., 2014b). The changes in terrigenous supply of fluvial discharge recorded by VC-04 core fit well with the decrease in the insolation during the late Holocene. We propose that the river runoff gradually decreased following the summer insolation. The findings of the present study will help in the characterisation of the evolutionary patterns of river-influenced continental shelf areas in Indian subcontinent.

6. Conclusions

The elemental distributions, sedimentation rates and grain size investigations performed on core sediment retrieved from the continental shelf adjacent to Rushikulya river mouth allowed us to identify paleoclimatic changes. The modern investigations of the surface sediments from the inner-mid continental shelf were conducted to understand the coastal dynamics and refine palaeo-environmental interpretations. The grain size distribution and elemental data of the surface sediments show that in the modern day condition, the elemental distribution such as TiO_2 and Al_2O_3 is largely controlled by the grain size of the sediments. The proportion of grain size (clay and silt vs sand) and elemental values (TiO_2 vs Al_2O_3) has been used to infer the changes in past terrigenous supply. Our results pointed out the period of higher river discharge from 6800 to 3100 cal BP followed by reduction of monsoonal strength. The drier conditions recorded off core sediments are corroborated with climatic changes recorded from other parts of Indian summer monsoon domain.

Acknowledgments

AA and PKM gratefully acknowledge the financial support provided by Inspire Fellowship from Department of Science and Technology. Authors also thank the X-ray facility at IISER Mohali for the help with XRD analyses. The authors confirm that they have no conflict of interest.

References

- Ameakwa, S., Kubota, K., Miyairi, Y., Seki, A., Kawakubo, Y., Sakai, S., Ajithprasad, P., Maemoku, H., Osada, T., Yokoyama, Y., 2016. Fossil otoliths, from the Gulf of Kutch, Western India, as a paleo-archive for the mid-to late-Holocene environment. *Quat. Int.* 397, 281–288.
- Anoop, A., Prasad, S., Krishnan, R., Naumann, R., Dulski, P., 2013a. Intensified monsoon and spatiotemporal changes in precipitation patterns in the NW Himalaya during the early-mid Holocene. *Quat. Int.* 313, 74–84.
- Anoop, A., Prasad, S., Plessen, B., Basavaiah, N., Gaye, B., Naumann, R., Menzel, P., Weise, S., Brauer, A., 2013b. Palaeoenvironmental implications of evaporative Gaylussite crystals from Lonar lake, Central India. *J. Quat. Sci.* 28, 349–359.
- Banerji, U.S., Pandey, S., Bhushan, R., Juyal, N., 2015. Mid-Holocene climate and land–sea interaction along the southern coast of Saurashtra, western India. *J. Asian Earth Sci.* 111, 428–439.
- Bhattacharya, S., Sengupta, R., 1994. Surface microtextures of heavy minerals from the Bay of Bengal off gopalpur, Orissa. *J. Geol. Soc. India* 44, 175–184.
- Boës, X., Rydberg, J., Martinez-Cortizas, A., Bindler, R., Renberg, I., 2011. Evaluation of conservative lithogenic elements (Ti, Zr, Al, and Rb) to study anthropogenic element enrichments in lake sediments. *J. Paleolimnol.* 46, 75–87.
- Briceno-Zuluaga, F.J., Sifeddine, A., Caquineau, S., Cardich, J., Salvatelli, R., Gutierrez, D., Ortlieb, L., Velasco, F., Boucher, H., Machado, C., 2016. Terrigenous material supply to the Peruvian central continental shelf (Pisco, 14° S) during the last 1000 years: paleoclimatic implications. *Clim. Past* 12, 787–798.
- Demske, D., Tarasov, P.E., Wünnemann, B., Riedel, F., 2009. Late glacial and Holocene vegetation, Indian monsoon and westerly circulation in the Trans-Himalaya recorded in the lacustrine pollen sequence from Tso Kar, Ladakh, NW India. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 279, 172–185.
- Dodia, R., Agrawal, D.P., Vora, A.B., 1984. New pollen data from Kashmir bogs: a summary. In: Whyte, R.O. (Ed.), *The Evolution of the East Asian Environment*, vol. 2. Hongkong Univ., Hongkong, pp. 569–578.
- Eglinton, T.I., Benitez-Nelson, B.C., Pearson, A., McNichol, A.P., Bauer, J.E., Druffel, E.R., 1997. Variability in radiocarbon ages of individual organic compounds from marine sediments. *Science* 277, 796–799.
- Enzel, Y., Ely, L.L., Mishra, S., Ramesh, R., Amit, R., Lazar, B., Rajaguru, S.N., Baker, V.R., Sandler, A., 1999. High-resolution Holocene environmental changes in the Thar Desert, northwestern India. *Science* 284, 125–128.
- Folk, R.L., 1980. *Petrology of Sedimentary Rocks*. Hemphill Publishing Company, Austin, TX, p. 182.
- Gingele, F.X., De Deckker, P., Hillenbrand, C.D., 2004. Late Quaternary terrigenous sediments from the Murray Canyons area, offshore South Australia and their implications for sea level change, palaeoclimate and palaeodrainage of the Murray–Darling Basin. *Mar. Geol.* 212, 183–197.
- González-Álvarez, R., Bernárdez, P., Pena, L.D., Francés, G., Prego, R., Diz, P., Vilas, F., 2005. Paleoclimatic evolution of the Galician continental shelf (NW of Spain) during the last 3000 years: from a storm regime to present conditions. *J. Mar. Syst.* 54, 245–260.
- Günther, F., Witt, R., Schouten, S., Mäusbacher, R., Daut, G., Zhu, L., Xu, B., Yao, T., Gleixner, G., 2015. Quaternary ecological responses and impacts of the Indian ocean summer monsoon at Nam Co, southern Tibetan plateau. *Quat. Sci. Rev.* 112, 66–77.
- Hameed, A., Achyuthan, H., Sekhar, B., 2006. Radiocarbon dates and Holocene sea level change along the Cuddalore and Odinur coast, Tamil Nadu. *Curr. Sci.* 91, 362–367.
- Hashimi, N.H., Nigam, R., Nair, R.R., Rajagopalan, G., 1995. Holocene sea level fluctuations on western Indian continental margin: an update. *J. Geol. Soc. India* 46, 157–162.
- Hjulstrom, F., 1935. *Studies of the Morphological Activities of Rivers as Illustrated by the River Fyris*, 25. *Bull. Geol. Inst., Univ., Uppsala*, pp. 221–527.
- Hjulstrom, F., 1939. Transportation of Detritus by Moving Water. In: Trask, P.D. (Ed.), *Recent Marine Sediments, a Symposium*. American Assn of Petroleum Geologists, Tulsa, Oklahoma, pp. 5–31.
- Hong, Y.T., Hong, B., Lin, Q.H., Shibata, Y., Hirota, M., Zhu, Y.X., Leng, X.T., Wang, Y., Wang, H., Yi, L., 2005. Inverse phase oscillations between the East Asian and Indian Ocean summer monsoons during the last 12000 years and paleo-El Niño. *Earth Planet. Sci. Lett.* 231, 337–346.
- Jain, S.K., Agarwal, P.K., Singh, V.P., 2007. *Hydrology and Water Resources of India*. Water Science and Technology Library. Springer-Verlag, Heidelberg, p. 1258.
- Juahir, H., Zain, S.M., Yusoff, M.K., Hanidza, T.T., Armi, A.M., Toriman, M.E., Mokhtar, M., 2011. Spatial water quality assessment of Langat River Basin (Malaysia) using environmetric techniques. *Environ. Monit. Assess.* 173, 625–641.
- Khandelwal, A., Gupta, H.P., 1999. Late Holocene climate and vegetation of Bhowania, Chilika lake, Orissa. *Gondwana Geol. Mag. Spec.* 4, 301–306.
- Khandelwal, A., Mohanti, M., García-Rodríguez, F., Scharf, B.W., 2008. Vegetation history and sea level variations during the last 13,500 years inferred from a pollen record at Chilika Lake, Orissa, India. *Veg. Hist. Archaeobotany* 17, 335–344.
- Kotlia, B.S., 2015. Updated quaternary climatic research in parts of the Third Pole. *Quat. Int.* 371, 1–4.
- Kotlia, B.S., Singh, A.K., Joshi, L.M., Dhaila, B.S., 2015. Precipitation variability in the Indian central Himalaya during last ca. 4,000 years inferred from a speleothem record: impact of Indian summer monsoon (ISM) and Westerlies. *Quat. Int.* 371, 244–253.
- Leipe, C., Demske, D., Tarasov, P.E., 2014. A Holocene pollen record from the northwestern Himalayan lake Tso Moriri: implication for palaeoclimatic and archaeological research. *Quat. Int.* 348, 93–112.
- Loska, K., Wiechula, D., 2003. Application of principal component analysis for the estimation of source of heavy metal contamination in surface sediments from the Rybnik Reservoir. *Chemosphere* 51, 723–733.
- Markose, V.J., Rajan, B., Kankara, R.S., Selvan, S.C., Dhanalakshmi, S., 2016. Quantitative analysis of temporal variations on shoreline change pattern along Ganjam district, Odisha, east coast of India. *Environ. Earth Sci.* 75, 1–17.
- Mascarenhas, A., 2004. Oceanographic validity of buffer zones for the east coast of India: a hydrometeorological perspective. *Curr. Sci.* 86, 399–406.
- McCave, I.N., Syvitski, J.P.M., 1991. *Principles and Methods of Particle Size Analysis*. In: Syvitski, J.P.M. (Ed.), *Principles, Methods, and Applications of Particle Size Analysis*. Cambridge University Press, New York, pp. 3–21.
- Mendes, I., Rosa, F., Dias, J.A., Schönfeld, J., Ferreira, O., Pinheiro, J., 2010. Inner shelf paleoenvironmental evolution as a function of land–ocean interactions in the vicinity of the Guadiana River, SW Iberia. *Quat. Int.* 221, 58–67.
- Miall, A.D., 2006. *The Geology of Fluvial Deposits: Sedimentary Facies, Basin Analysis and Petroleum Geology*. Springer-Verlag, Berlin.
- Milliman, J.D., Emery, K.O., 1968. Sea levels during the past 35,000 years. *Science* 162, 1121–1123.
- Minyuk, P.S., Borkhodoev, V.Y., Wennrich, V., 2013. Inorganic data from El'gygytyn Lake sediments: stages 6–11. *Clim. Past. Discuss.* 9, 393–433.
- Mishra, P., Mohanty, P.K., Murty, A.S.N., Sugimoto, T., 2001. Beach profile studies near an artificial open-coast port along south Orissa, east coast of India. *J. Coast. Res.* 164–171.
- Mishra, P., Praveen, K., Anoop, A., Schettler, G., Prasad, S., Jehangir, A., Menzel, P., Naumann, R., Yousuf, A.R., Basavaiah, N., Deenadayalan, K., Wiesner, M.G., Gaye, B., 2015. Reconstructed late quaternary hydrological changes from Tso Moriri lake, NW Himalaya. *Quat. Int.* 371, 76–86.
- Morrison, D., 1967. *Multivariate Statistical Methods*. McGrawHill, New York.
- Nizou, J., Hanebuth, T.J., Heslop, D., Schwenk, T., Palamenghi, L., Stuut, J.B., Henrich, R., 2010. The Senegal River mud belt: a high-resolution archive of paleoclimatic change and coastal evolution. *Mar. Geol.* 278, 150–164.
- Panigrahy, P.K., Das, J., Das, S.N., Sahoo, R.K., 1999. Evaluation of the influence of various physico-chemical parameters on coastal water quality, around Orissa, by factor analysis. *Indian J. Mar. Sci.* 28, 360–364.
- Perez, L., García-Rodríguez, F., Hanebuth, T.J., 2016. Variability in terrigenous

- sediment supply offshore of the Rio de la Plata (Uruguay) recording the continental climatic history over the past 1200 years. *Clim. Past* 12, 623–634.
- Phadtare, N.R., 2000. Sharp decrease in summer monsoon strength 4000–3500 cal yr BP in the Central Higher Himalaya of India based on pollen evidence from alpine peat. *Quat. Res.* 53, 122–129.
- Ponton, C., Giosan, L., Eglinton, T.I., Fuller, D.Q., Johnson, J.E., Kumar, P., Collett, T.S., 2012. Holocene aridification of India. *Geophys. Res. Lett.* 39, 1–6.
- Prasad, S., Enzel, Y., 2006. Holocene paleoclimates of India. *Quat. Res.* 66, 442–453.
- Prasad, S., Kusumgar, S., Gupta, S.K., 1997. A mid to late Holocene record of palaeoclimatic changes from Nal Sarovar: a palaeodesert margin lake in western India. *J. Quat. Sci.* 12, 153–159.
- Prasad, V., Farooqui, A., Sharma, A., Phartiyal, B., Chakraborty, S., Bhandari, S., Raj, R., Singh, A., 2014a. Mid–late Holocene monsoonal variations from mainland Gujarat, India: a multi-proxy study for evaluating climate culture relationship. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 397, 38–51.
- Prasad, S., Anoop, A., Riedel, N., Sarkar, S., Menzel, P., Basavaiah, N., Krishnan, R., Fuller, D., Plessen, B., Gaye, B., Röhl, U., 2014b. Prolonged monsoon droughts and links to Indo-Pacific warm pool: a Holocene record from Lonar Lake, central India. *Earth Planet. Sci. Lett.* 391, 171–182.
- Prins, M.A., Postma, G., Cleveringa, J., Cramp, A., Kenyon, N.H., 2000. Controls on terrigenous sediment supply to the Arabian sea during the late quaternary: the Indus fan. *Mar. Geol.* 169, 327–349.
- Raj, R., Chamyal, L.S., Prasad, V., Sharma, A., Tripathi, J.K., Verma, P., 2015. Holocene climatic fluctuations in the Gujarat Alluvial Plains based on a multiproxy study of the Pariyaj Lake archive, western India. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 421, 60–74.
- Ramakrishnan, M., Nanda, J.K., Augustine, P.F., 1998. Geological evolution of the Proterozoic eastern ghats mobile belt. *Geol. Surv. India Spec. Publ.* 44, 1–21.
- Ramsey, C.B., 2001. Development of the radiocarbon calibration program. *Radiocarbon* 43, 355–364.
- Ramsey, C.B., 2009. Bayesian analysis of radiocarbon dates. *Radiocarbon* 51, 337–360.
- Rao, C.B., 1957. Beach erosion and concentration of heavy mineral sands. *J. Sediment. Res.* 27, 143–147.
- Rao, N.S., Misra, S., 2009. Sources of monazite sand in southern Orissa beach placer, eastern India. *J. Geol. Soc. India* 74, 357–362.
- Rao, R.G., Sahoo, P., Panda, N.K., 2001. Heavy mineral sand deposit of Orissa. *Explor. Res. Atomic Minerals* 13, 23–52.
- Rawat, S., Gupta, A.K., Sangode, S.J., Srivastava, P., Nainwal, H.C., 2015. Late Pleistocene–Holocene vegetation and Indian summer monsoon record from the Lahaul, Northwest Himalaya, India. *Quat. Sci. Rev.* 114, 167–181.
- Reid, M.K., Spencer, K.L., 2009. Use of principal components analysis (PCA) on estuarine sediment datasets: The effect of data pre-treatment. *Environ. Pollut.* 157, 2275–2281.
- Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Bronk Ramsey, C., Buck, C.E., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., 2013. IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon* 55, 1869–1887.
- Rosa, F., Dias, J.A., Mendes, I., Ferreira, Ó., 2011. Mid to late Holocene constraints for continental shelf mud deposition in association with river input: the Guadiana Mud Patch (SW Iberia). *Geo-Marine Lett.* 31, 109–121.
- Sappa, G., Ergul, S., Ferranti, F., 2014. Geochemical Modelling and Multivariate Statistical Evaluation of Trace Elements in Arsenic Contaminated Groundwater Systems of Viterbo Area (Central Italy), vol. 3. SpringerPlus, p. 237.
- Sinha, A., Berkelhammer, M., Stott, L., Mudelsee, M., Cheng, H., Biswas, J., 2011. The leading mode of Indian Summer Monsoon precipitation variability during the last millennium. *Geophys. Res. Lett.* 38, L15703.
- Trivedi, A., Chauhan, M.S., 2009. Holocene vegetation and climate fluctuations in Northwest Himalaya based on pollen evidence from Surinsar Lake, Jammu region, India. *J. Geol. Soc. India* 74, 402–412.
- Tu, L.Y., Zhou, X., Cheng, W.H., Liu, X.Y., Yang, W.Q., Wang, Y.H., 2016. Holocene East Asian winter monsoon changes reconstructed by sensitive grain size of sediments from Chinese coastal seas: a review. *Quat. Int.* 44 (PA), 82–90. <http://dx.doi.org/10.1016/j.quaint.2016.03.024>.
- Varghese, S., Roychaudhuri, S., Mashood, P.A., Tripathi, S.K., 2015. Geomorphology of the offshore of Rushikulya river mouth in northern Bay of Bengal. *Indian J. Geosci.* 69, 233–240.
- Visher, G.S., 1969. Grain size distributions and depositional processes. *J. Sediment. Res.* 39, 1074–1106.
- Wang, K., Zheng, H., Tada, R., Irino, T., Zheng, Y., Saito, K., Karasuda, A., 2014. Millennial-scale East Asian Summer Monsoon variability recorded in grain size and provenance of mud belt sediments on the inner shelf of the East China Sea during mid-to late Holocene. *Quat. Int.* 349, 79–89.
- Wasson, R.J., Smith, G.I., Agrawal, D.P., 1984. Late quaternary sediments, minerals, and inferred geochemical history of Didwana Lake, Thar Desert, India. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 46, 345–372.
- Wünnemann, B., Demske, D., Tarasov, P., Kotlia, B.S., Reinhardt, C., Bloemendal, J., Diekmann, B., Hartmann, K., Krois, J., Riedel, F., Arya, N., 2010. Hydrological evolution during the last 15kyr in the Tso Kar lake basin (Ladakh, India), derived from geomorphological, sedimentological and palynological records. *Quat. Sci. Rev.* 29, 1138–1155.
- Yoo, D.G., Kim, S.P., Chang, T.S., Kong, G.S., Kang, N.K., Kwon, Y.K., Nam, S.L., Park, S.C., 2014. Late Quaternary inner shelf deposits in response to late Pleistocene–Holocene sea level changes: Nakdong River, SE Korea. *Quat. Int.* 344, 156–169.
- Zhang, J., Liu, C.L., 2002. Riverine composition and estuarine geochemistry of particulate metals in China—weathering features, anthropogenic impact and chemical fluxes. *Estuar. Coast. Shelf Sci.* 54, 1051–1070.
- Zhao, B., Wang, Z., Chen, J., Chen, Z., 2008. Marine sediment records and relative sea level change during late Pleistocene in the Changjiang delta area and adjacent continental shelf. *Quat. Int.* 186, 164–172.
- Zheng, Y., Anderson, R.F., Froelich, P.N., Beck, W., McNichol, A.P., Guilderson, T., 2002. Challenges in radiocarbon dating organic carbon in opal-rich marine sediments. *Radiocarbon* 44, 123–136.
- Zheng, Y., Kissel, C., Zheng, H.B., Laj, C., Wang, K., 2010. Sedimentation on the inner shelf of the East China Sea: magnetic properties, diagenesis and paleoclimate implications. *Mar. Geol.* 268, 34–42.